

Dogs: Flea and Tick Products

Oral/Systemic Products			
	NAME (COMPANY)		
1	Bravecto (Intervet)	ACTIVE INGREDIENT	Fluralaner
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs only (6 months and over 4.4 pounds)
		FREQUENCY	Lone star ticks: 8 weeks; fleas, other ticks: 12 weeks
		EFFECTS	Kills fleas; kills lone star tick, black-legged (deer) tick, American dog tick, brown dog tick
		COMMENTS	Administer with food. May be used with other medications such as vaccines, deworming medications, antibiotics, and corticosteroids
	NAME (COMPANY)		
2	Capstar (Elanco)	ACTIVE INGREDIENT	Nitenpyram
		TYPE	Tablet
		PET, MINIMUM AGE	Dogs (4 weeks and over 2 pounds) Cats (4 weeks and over 2 pounds)
		FREQUENCY	As needed, up to once per day
		EFFECTS	Kills fleas
		COMMENTS	Occasionally, dog or cat will start scratching itself as fleas begin to die. The scratching behavior is temporary and is a reaction to the fleas, not the drug. In cats and dogs, this may appear as temporary signs of hyperactivity, panting, vocalization and excessive grooming/licking. May be used together with other products, including heartworm preventives, corticosteroids, antibiotics, vaccines, de-worming medications, shampoos and other flea products



	NAME (COMPANY)		
3	Comfortis (Elanco)/ Acuguard (VCA)	ACTIVE INGREDIENT	Spinosad
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (14 weeks and over 3.3 pounds) Cats (14 weeks and over 2 pounds)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas
		COMMENTS	Administer with food. May be used with other frequently used veterinary products such as vaccines, deworming medications, antibiotics, corticosteroids, shampoos, and other flea products
	NAME (COMPANY)		
4	Nexgard (Merial)	ACTIVE INGREDIENT	Afoxolaner
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (8 weeks and over 4 pounds)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas; black-legged (deer) tick, American dog tick, lone star tick, brown dog tick
		COMMENTS	Can be administered with or without food
	NAME (COMPANY)		
5	Program (Elanco)	ACTIVE INGREDIENT	Lufenuron
		TYPE	Oral suspension, Chewable tablet
		PET, MINIMUM AGE	Dogs (over 4 weeks), Cats (over 6 weeks)

		FREQUENCY	Monthly
		EFFECTS	Breaks flea life cycle
		COMMENTS	Does not kill adult fleas; breaks the flea life cycle at the egg stage to prevent eggs from developing into adults. Pre-existing immature fleas in the environment may continue to develop and emerge after treatment. Concurrent use of adulticidal insecticides may be used to control adult fleas. Administer with food
	NAME (COMPANY)		
6	Sentinel (Virbac)	ACTIVE INGREDIENT	Lufenuron, Milbemycin
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (over 4 weeks and over 2 pounds)
		FREQUENCY	Monthly
		EFFECTS	Breaks flea life cycle; prevents heartworm; controls hookworms, roundworms and whipworms
		COMMENTS	Does not kill adult fleas; breaks the flea life cycle at the egg stage to prevent eggs from developing into adults. Pre-existing immature fleas in the environment may continue to develop and emerge after treatment. Concurrent use of adulticidal insecticides may be used to control adult fleas.
	NAME (COMPANY)		
7	Sentinel Spectrum (Virbac)	ACTIVE INGREDIENT	Lufenuron, Milbemycin, Praziquantel
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (over 6 months and over 2.8 pounds)
		FREQUENCY	Monthly
		EFFECTS	Breaks flea life cycle; prevents heartworm; controls hookworms, roundworms, whipworms, tapeworms

		COMMENTS	Does not kill adult fleas; breaks the flea life cycle at the egg stage to prevent eggs from developing into adults. Pre-existing immature fleas in the environment may continue to develop and emerge after treatment. Concurrent use of adulticidal insecticides may be used to control adult fleas.
	NAME (COMPANY)		
8	Simparica (Zoetis)	ACTIVE INGREDIENT	Sarolaner
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (over 6 months and over 2.8 pounds)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas; kills lone star tick, Gulf coast tick, American dog tick, brown dog tick
		COMMENTS	May be administered with or without food
	NAME (COMPANY)		
9	Trifexis (Elanco)	ACTIVE INGREDIENT	Spinosad, Milbemycin
		TYPE	Chewable tablet
		PET, MINIMUM AGE	Dogs (over 8 weeks and over 5 pounds)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas. Prevents heartworm when given 3 consecutive monthly doses. Kills adult hookworms, roundworms, whipworms
		COMMENTS	Puppies less than 14 weeks of age may experience vomiting. Administer with food
Topical Products			
	NAME (COMPANY)		

10	Activyl Tick PLUS (Merck)	ACTIVE INGREDIENT	Indoxacarb, Permethrin
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 8 weeks and over 4 pounds)
		FREQUENCY	30 days
		EFFECTS	Kills fleas; American dog tick, brown dog tick, deer tick, lone star tick
		COMMENTS	Waterproof. Consult a veterinarian before using this product on debilitated, aged, and medicated animals. Do not use on pregnant or nursing animals. Harmful or fatal if used on cats.
	NAME (COMPANY)		
11	Advantage (Bayer)	ACTIVE INGREDIENT	Imidacloprid
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 7 weeks) Cats (over 8 weeks)
		FREQUENCY	Monthly; In case of severe flea infestation, retreatment may be necessary earlier than 4 weeks. Do not retreat more often than once every 7 days.
		EFFECTS	Kills fleas
		COMMENTS	Waterproof. Consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medications
	NAME (COMPANY)		
12	Advantage II (Bayer)	ACTIVE INGREDIENT	Imidacloprid, Pyriproxyfen
		TYPE	Spot-on



		PET, MINIMUM AGE	Dogs (over 7 weeks), Cats (over 8 weeks)
		FREQUENCY	Monthly; In case of severe flea infestation, retreatment may be necessary earlier than 4 weeks. Do not retreat more often than once every 7 days.
		EFFECTS	Kills fleas
		COMMENTS	Waterproof. Consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medications
	NAME (COMPANY)		
13	Advantage Multi (Bayer)	ACTIVE INGREDIENT	Imidacloprid, Moxidectin
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 7 weeks and over 3 pounds), Cats (over 9 weeks and over 2 pounds), Ferrets (cat product only, for ferrets over 2 pounds)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas. Prevents heartworms and (except in ferrets) controls hookworms, roundworms, whipworms
		COMMENTS	Do not allow pet to lick application site; ingestion may result in serious adverse reactions. Shampooing or water immersion 4 days after application does not reduce flea control effectiveness, but shampooing as often as once weekly may reduce flea control effectiveness.
	NAME (COMPANY)		
14	Certifect (Merial)	ACTIVE INGREDIENT	Fipronil, Methoprene, Amitraz
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 8 weeks)
		FREQUENCY	Monthly (ticks)



		EFFECTS	Kills fleas; kills deer ticks, Gulf coast ticks, black legged ticks, brown dog ticks, American dog ticks, lone star ticks, yellow dog ticks, European dog tick, wood ticks, and paralysis ticks; starts killing ticks within 6 hours; aids in control of sarcoptic mange infestations; eliminates chewing lice infestations
		COMMENTS	Waterproof. Can be used on breeding, pregnant and lactating bitches; Consult a veterinarian before using on dogs that are medicated, debilitated, aged or obese.
	NAME (COMPANY)		
15	Effitix (Virbac)	ACTIVE INGREDIENT	Fipronil, Permethrin
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 12 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas. Kills brown dog ticks, American dog ticks, lone star ticks, deer ticks. Kills mosquitoes. Prevents blood-feeding. Kills lice. Aids in control of sarcoptic mites.
		COMMENTS	Waterproof. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs, or animals known to be sensitive to pesticide products.
	NAME (COMPANY)		
16	Frontline Plus (Merial)	ACTIVE INGREDIENT	Fipronil, Methoprene
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 6 weeks) Cats (over 8 weeks)
		FREQUENCY	Monthly (ticks, mosquitoes, flies, lice)
		EFFECTS	Kills fleas; kills deer ticks, brown dog ticks, American dog ticks, lone star ticks. Eliminates chewing lice. Aids in control of sarcoptic mites



		COMMENTS	Waterproof. Can be used on breeding, pregnant and lactating dogs and cats.
	NAME (COMPANY)		
17	Frontline Tritak for Dogs (Merial)	ACTIVE INGREDIENT	Fipronil, Cyphenothrin, Methoprene
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 12 weeks)
		FREQUENCY	30 days
		EFFECTS	Kills fleas; American ticks, brown dog ticks, lone star ticks, deer ticks; chewing lice. Controls sarcoptic mites.
		COMMENTS	Waterproof. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction.
	NAME (COMPANY)		
18	K9 Advantix (Bayer)	ACTIVE INGREDIENT	Imidacloprid, Permethrin
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 7 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas; kills deer ticks, American dog ticks, brown dog ticks and lone star ticks. Kills mosquitoes. Prevents blood-feeding by biting flies. Kills lice.
		COMMENTS	Waterproof. Consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medications. Harmful or fatal if applied to cats or if cats groom recently treated dogs.
	NAME (COMPANY)		



19	K9 Advantix II (Bayer)	ACTIVE INGREDIENT	Imidacloprid, Permethrin, Pyriproxyfen
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 7 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas; kills deer ticks, American dog ticks, brown dog ticks, lone star ticks. Kills mosquitoes. Kills lice. Prevents blood-feeding by biting flies.
		COMMENTS	Waterproof. Consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medications. Harmful or fatal if applied to cats or if cats groom recently treated dogs.
	NAME (COMPANY)		
20	Parastar (Novartis)	ACTIVE INGREDIENT	Fipronil
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (over 8 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas; Kills deer ticks, brown dog ticks, American dog ticks, and lone star ticks. Kills chewing lice. Controls sarcoptic mites.
		COMMENTS	Waterproof. Can be used on breeding, pregnant, or lactating dogs. Consult a veterinarian before using this product on debilitated, aged, or medicated dogs or on dogs with known organ dysfunction.
	NAME (COMPANY)		
21	Parastar PLUS (Novartis)	ACTIVE INGREDIENT	Fipronil, Cyphenothrin
		TYPE	Spot-on



		PET, MINIMUM AGE	Dogs (over 12 weeks)
		FREQUENCY	4 weeks
		EFFECTS	Kills fleas. Kills deer ticks, brown dog ticks, American dog ticks, and lone star ticks. Kills chewing lice. Controls sarcoptic mites.
		COMMENTS	Waterproof. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs or on dogs with known organ dysfunction. Harmful or fatal if applied to cats or if cats groom recently treated dogs.
	NAME (COMPANY)		
22	Preventic (Virbac)	ACTIVE INGREDIENT	Amitraz
		TYPE	Collar
		PET, MINIMUM AGE	Dogs (over 12 weeks)
		FREQUENCY	90 days
		EFFECTS	Kills ticks
		COMMENTS	Water resistant. Does not kill fleas. Consult a veterinarian before using on debilitated, aged, medicated, pregnant or nursing animals. Do not apply to cats. Do not allow dogs to ingest collar or parts of collar; contact veterinarian immediately if ingestion of collar is suspected.
	NAME (COMPANY)		
23	Revolution (Zoetis)	ACTIVE INGREDIENT	Selamectin
		TYPE	Spot-on
		PET, MINIMUM AGE	Dogs (6 weeks) Cats (8 weeks)
		FREQUENCY	Monthly



		EFFECTS	Kills fleas, ear mites; heartworm preventive. Dogs: Controls sarcoptic mites. Kills American dog ticks. Cats: Treats and controls roundworm and hookworms.
		COMMENTS	Do not use in sick, debilitated or underweight animal
	NAME (COMPANY)		
24	Scalibor (Merck)	ACTIVE INGREDIENT	Deltamethrin
		TYPE	Collar
		PET, MINIMUM AGE	Dogs only (12 weeks)
		FREQUENCY	6 months
		EFFECTS	Kills fleas; kills brown dog tick, American dog tick, deer ticks.
		COMMENTS	Water resistant. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing dogs. Use on dogs only.
	NAME (COMPANY)		
25	Seresto (Bayer)	ACTIVE INGREDIENT	Flumethrin, Imidacloprid
		TYPE	Collar
		PET, MINIMUM AGE	Dogs (over 7 weeks), Cats (over 10 weeks)
		FREQUENCY	8 months
		EFFECTS	Kills fleas; kills deer ticks, American dog ticks, brown dog ticks, lone star ticks. Kills chewing lice. Controls sarcoptic mites. Cats: Remove ticks already on cat at time of collar application as they may not be killed immediately



		COMMENTS	Water resistant; for animals that are bathed or swim more than once per month, control duration is reduced to 5 months. Consult veterinarian before using this product on debilitated, aged, breeding, pregnant, or nursing animals.
	NAME (COMPANY)		Dinotefuran, Pyriproxyfen
26	Vectra (Ceva)	ACTIVE INGREDIENT	Spot-on
		TYPE	
		PET, MINIMUM AGE	Dogs (over 8 weeks), Cats (over 8 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas
		COMMENTS	Waterproof for one month. Consult veterinarian before using this product on debilitated, aged, breeding, pregnant, or nursing animals or on animals with known sensitivity to pesticides.
	NAME (COMPANY)		Dinotefuran, Pyriproxyfen, Permethrin
27	Vectra 3D (Ceva)	ACTIVE INGREDIENT	Spot-on
		TYPE	
		PET, MINIMUM AGE	Dogs (over 7 weeks)
		FREQUENCY	Monthly
		EFFECTS	Kills fleas within 6 hours. Kills American dog ticks, lone star ticks, brown dog ticks, deer ticks. Kills mosquitoes, lice, sand flies, mites
		COMMENTS	Water resistant. Consult veterinarian before using this product on debilitated, aged, breeding, pregnant, or nursing animals or on animals with known sensitivity to pesticides. Harmful or fatal if applied to cats or if cats groom recently treated dogs.

